

COMPOSITION

Each film coated tablet contains Thiamine Mononitrate USP (Vitamin B_1) 100 mg, Pyridoxine Hydrochloride BP (Vitamin B_6) 200 mg & Cyanocobalamin BP (Vitamin B_{12}) 200 mcg.

PHARMACOLOGY

 ${\bf V3N}^{\rm @}$ is a combined preparation of Vitamin B1 (Thiamine Mononitrate), Vitamin B6 (Pyridoxine Hydrochloride) and Vitamin B12 (Cyanocobalamin). These vitamins play an essential role as co-enzyme in the metabolism of nervous system. ${\bf V3N}^{\rm @}$ supports the regeneration of nerve fibers and myelin sheath by activation of the nerve cell metabolism and the natural repair mechanism.

INDICATION

Chronic pains like sciatica, lumbago, trigeminal neuralgia, facial paralysis, optic neuritis as well as neuropathy & cardiac complications. Besides these, this combination may be used in the deficiency of these vitamins.

DOSAGE AND ADMINISTRATION

1 to 3 tablets per day or as directed by the physician.

CONTRAINDICATION

It should not be used in patients with known hypersensitivity to Thiamine, Cobalt or any other component of these preparations.

WARNING AND PRECAUTION

It should be avoided in case of hypersensitivity to Thiamine, Cobalt or any other component of these preparations. Cyanocobalamin should not be given before a diagnosis has been fully established because of the masking symptoms of subacute degeneration of the spinal cord.

SIDE EFFECTS

Very rarely hypersensitive reaction and sensitivity disturbance may occur.

USE IN PREGNANCY AND LACTATION

No risks have become known associated with the use of $V3N^{\circ}$ during pregnancy at the recommended dosage. Vitamin B₁, B₆ & B₁₂ are secreted into the breast milk but risks of overdose for the infant are not known.

USE IN CHILDREN

Vitamin B₁ & B₆ are not recommended under 3 years.

DRUG INTERACTION

Vitamin B6 (Pyridoxine Hydrochloride) reduces therapeutic effect of Levodopa but it does not occur if a dopa decarboxylase inhibitor is also given. Neomycin, Chloramphenicol, Aminosalicylic Acid & H₂ receptor antagonist may reduce Vitamin B₁₂ absorption.

OVERDOSAGE

Excess Vitamin B is readily excreted, so no serious problems occur.

STORAGE

Store below 30°C temperature in a cool and dry place. Protect from light and moisture. Keep out of the reach of children.

HOW SUPPLIED

Each box contains 50 tablets in Alu-PVDC blister pack.



Manufactured by:

NIPRO JMI Pharma Ltd.

Chauddagram, Cumilla, Bangladesh.