Visnil

Tiemonium Methylsulphate INN

COMPOSITION

Visnil Tablet: Each film coated tablet contains Tiemonium Methylsulphate INN 50 mg.
Visnil Syrup: Each 5 ml syrup contains Tiemonium Methylsulphate INN 10 mg.
Visnil Injection: Each 2 ml ampoule contains sterile solution of Tiemonium Methylsulphate INN 5 mg.

PHARMACOLOGY

Visnil (Tiemonium Methylsulphate) is an antispasmodic drug that reduces muscles spasm of the intestine, biliary system, bladder and uterus.

INDICATIONS

Tiemonium Methylsulphate is an antispasmodic drug. It reduces muscle spasms of the intestine, biliary system, uterus and urinary bladder. It is indicated for the pain in gastrointestinal, biliary, urinary and gynecological disease such as gastroenteritis, diarrhoea, dysentery, biliary colic, enterocolitis, cholecystitis, colonopathy, mild cystitis and spasmodic dysmenorrhoea.

DOSAGE AND ADMINISTRATION

Adult: Usual dose is 2-6 tablets or 3-9 teaspoonfuls syrup daily in divided doses.
Children: 3 ml/kg or 6 mg/kg body weight daily in divided doses.
Dosage of Injection: 1 ampoule of Tiemonium Methylsulphate Injection 3 times daily, through the Intravenous or Intramuscular route slowly.

CONTRAINDICATIONS

It should not be used in urethroprostatic disorder involving a risk of urine retention. It is contraindicated in patient with having risk of angle closure glaucoma.

WARNING AND PRECAUTION

Caution should be taken during treatment of patients with disorders of the prostate. Caution should also be taken in case of chronic bronchitis, coronary insufficiency, ambient hyperthermia, renal and hepatic insufficiency. The risks of visual disturbances can make it dangerous to drive or use machines.

SIDE EFFECTS

Tiemonium Methylsulphate may have the risk of hypotension and tachycardia in certain individuals.

USE IN PREGNANCY AND LACTATION

The results of animal studies of Tiemonium Methylsulphate did not reveal any teratogenic effects; no deformities have been reported untill now with normal use. In absence of sufficient data, prudence should be the rule for nursing mothers although no problems have been reported with normal use.

USE IN CHILDREN AND ADOLESCENTS

Safety and effectiveness of Tiemonium Methylsulphate in paediatric patients have not been established.

DRUG INTERACTIONS

It should not be used with other drugs. Without prior consultation with a registered physician avoid possible drug interaction.

OVERDOSAGE

The overdosage may appear by a tachycardia, an agitation, respiratory disorders. In case of overdosage immediately consult with the attending practitioners.

STORAGE

Visnil Tablet: Store below 30°C temperature in a cool and dry place. Protect from light and moisture. Keep out of the reach of children.
Visnil Syrup and Injection: Store below 30°C temperature in a cool and dry place. Protect from light. Keep out of the reach of children.

HOW SUPPLIED

Visnil Tablet: Each box contains 50 tablets in Alu-PVC blister pack.
Visnil Syrup: Each Amber PET bottle contains 50 ml & 100 ml syrup.
Visnil Injection: Each box contains 5 ampoules in Alu-PVC blister pack.

14020000039/02

NIPRO JMI Pharma

Manufactured by:
NIPRO JMI Pharma Ltd.
Chauddagram, Cumilla, Bangladesh.